

Patient's Name:

Allergies:

Medication Instruction Card

Bring This Card To All Clinic Appointments And Hospital Admissions

***** Please call 584-7001 for refills 7 days in advance of clinic appointment or before refill is needed *****

Immunosuppression	Strength	Directions	Morning	Midmorning	Afternoon	Evening	Bedtime
Cellcept (mycophenolate)							
Neoral (cyclosporine) Prograf (tacrolimus)		Take medication every 12 hours as directed. Remember to hold this medication until after blood work is drawn then take this medication.					
Rapamune (sirolimus)							
Deltasone (prednisone)		Take with food as directed.					
Other Medications	Strength	Directions	Morning	Midmorning	Afternoon	Evening	Bedtime
Bactrim DS / SS Pentamidine Treatment <i>To Prevent Bacterial Infection</i>							
Valtrex (valacyclovir) Valcyte (valganciclovir)* <i>To Prevent Viral Infection</i>							
Diflucan (fluconazole) Nystatin suspension <i>To Prevent Fungal Infection</i>							
Lasix (furosemide) Demadex (torsemide)							
Pepcid (famotidine) Protonix (pantoprazole) Prevacid (lansoprazole) Prilosec (omeprazole) Nexium (esomeprazole) <i>To Suppress Stomach Acid</i>							

**Patients taking a half tablet of Valcyte (valganciclovir) should cut the tablet themselves. All others wear gloves when handling tablets.*

